## Myosin Light Chain 2 (Phospho-Ser19) Antibody

Catalog No: #11114

Package Size: #11114-1 50ul #11114-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description	
Product Name	Myosin Light Chain 2 (Phospho-Ser19) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Rt
Specificity	The antibody detects endogenous level of Myosin Light Chain 2 only when phosphorylated at serine 19.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 19 (A-T-S(p)-N-V) derived from Human Myosin Light
	Chain 2.
Target Name	Myosin Light Chain 2
Modification	Phospho
Other Names	MLC-2a; MLC2a; MLRA; MYL7; MYLC2A
Accession No.	Swiss-Prot: P19105NCBI Protein: NP_006088.2
Uniprot	P19105
GeneID	10627;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details		
Predicted MW: 18kd		
Western blotting: 1:500~1:1000		
Immunohistochemistry: 1:50~1:100		

Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Myosin Light Chain 2 (Phospho-Ser19) Antibody #11114 (left) or the same antibody preincubated with blocking peptide (right).



Western blot analysis of extracts from 293 cells untreated or treated with starvation using Myosin Light Chain 2 (Phospho-Ser19) Antibody #11114.

## Background

Myosin regulatory subunit that plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis, receptor capping, and cell locomotion

Janiak A, et al. (2006) Mol Biol Cell. Apr; 17(4): 1606-1619.

Croft DR, et al. (2006) Mol Cell Biol. 2 Jun; 26(12): 4612-4627

Li Z, et al. (2006) Mol Cell Biol. Jun; 26(11): 4240-4256

Note: This product is for in vitro research use only