#### **Product Datasheet**

# HER2 (Phospho-Tyr1112) Conjugated Antibody

Catalog No: #C11903



Package Size: #C11903-AF350 100ul #C11903-AF405 100ul #C11903-AF488 100ul #C11903-AF555 100ul #C11903-AF594 100ul #C11903-AF647 100ul #C11903-AF680 100ul #C11903-AF750 100ul #C11903-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	HER2 (Phospho-Tyr1112) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of HER2 only when phosphorylated at tyrosine 1112.
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 1112 (Q-R-Y(p)-S-E) derived from Human HER2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	C-erbB-2;NGL;NEU proto-oncogene;MLN 19
Accession No.	Swiss-Prot#:P04626NCBI Gene ID:2064NCBI mRNA#:NM_001005862.2NCBI Protein#: NP_001005862.1
Uniprot	P04626
GenelD	2064;
Excitation Emission	AF350: 346nm/442nm
	AF405: 401nm/421nm
	AF488: 493nm/519nm
	AF555: 555nm/565nm
	AF594: 591nm/614nm
	AF647: 651nm/667nm
	AF680: 679nm/702nm
	AF750: 749nm/775nm
Calculated MW	150
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°Cin dark for 6 months

### **Application Details**

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250 AF405 conjugated: most applications: 1: 50 - 1: 250 AF488 conjugated: most applications: 1: 50 - 1: 250 AF555 conjugated: most applications: 1: 50 - 1: 250 AF594 conjugated: most applications: 1: 50 - 1: 250 AF647 conjugated: most applications: 1: 50 - 1: 250 AF680 conjugated: most applications: 1: 50 - 1: 250 AF750 conjugated: most applications: 1: 50 - 1: 250 Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

## **Product Description**

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

#### Background

Essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. GP30 is a potential ligand for this receptor. Not activated by EGF, TGF-alpha and amphiregulin.

Note: This product is for in vitro research use only