

Recombinant Human IL-8 (72a.a.)(rHu IL-8/CXCL8)

Catalog No: #701081

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Recombinant Human IL-8 (72a.a.)(rHu IL-8/CXCL8)
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	> 97 % by SDS-PAGE and HPLC analyses.
Species Reactivity	Hu
Target Name	rHu IL-8 CXCL8
Other Names	(Ser-IL-8)72, GCP IL-8 protein I, IL8 NAP1 form III, LYNAF, MDNCF-c, NAF
Accession No.	accession:P10145 GeneID:3576
Uniprot	P10145
GeneID	3576;
Calculated MW	Approximately 8.4 kDa, a singl
SDS-PAGE MW	Sterile Filtered White lyophil
Target Sequence	SAKELRCQCI KTYSKPFHPK FIKELRVIES GPHCANTEII VKLSDGRELCLDPKENWVQR VVEKFLKRAE NS
Formulation	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Storage	This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze thaw cycles.

Background

Interleukin-8 (IL-8) is encoded by the IL8 gene and produced by macrophages and other cell types such as epithelial cells. It is also synthesized by endothelial cells, which store IL-8 in their storage vesicles. There are many receptors capable to bind IL-8, the most affinity to IL-8 are receptors CXCR1, and CXCR2. As a member of the CXC chemokine family, function of IL-8 is the induction of chemotaxis in its target cells, like neutrophil granulocytes, basophils, and T-cells. IL-8 (72a.a.) has a 5-10-fold higher activity on neutrophil activation, compared to IL-8 (77a.a.). IL-8 is often associated with inflammation and has been cited as a proinflammatory mediator in gingivitis and psoriasis.

References

1. Modi WS, Dean M, Seuanez HN, et al. 1990. Hum Genet. 84:185-7.
2. Wolff B, Burns AR, Middleton J, et al. 1998. J Exp Med. 188:1757-62.
3. Utgaard JO, Jahnsen FL, Bakka A, et al. 1998. J Exp Med. 188:1751-6.
4. Van Damme J, Rampart M, Conings R, et al. 1990. Eur J Immunol. 20:2113-8.

Note: This product is for in vitro research use only