## Histone H3K27me2 Polyclonal Antibody

Catalog No: #HW007

Signalway Antibody



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size: #HW007-1 50ul #HW007-2 100ul


<b>-</b>			
1114	3 C C I	IAH	n
ᆫ	escri	ĮΟU	ווטו

Becomplien		
Product Name	Histone H3K27me2 Polyclonal Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Isotype	IgG	
Purification	Affinity purification	
Applications	WB,IF	
Species Reactivity	Human,Mouse,Rat	
Immunogen Type	Peptide	
Immunogen Description	A synthetic methylated peptide of human histone H3	
Target Name	Histone H3	
Modification	Methyl	
Other Names	H3.4;H3/g;H3FT;H3t;HIST3H3;Histone H3;HIST1H3A	
Accession No.	Uniprot:Q16695GeneID:8290	
Uniprot	Q16695	
GeneID	8290	
SDS-PAGE MW	17KDa	
Concentration	1.0mg/ml	
Formulation	PBS with 0.02% sodium azide,50% glycerol,pH7.3.	
Storage	Store at -20°C. Avoid freeze / thaw cycles.	

## Application Details

WB 1:500 - 1:2000IF 1:50 - 1:200

## **Images**



Western blot analysis of extracts of various cell lines, using DiMethyl-Histone H3-K27 antibody.

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Note: This product is for in vitro research use only