


Histone H4R3me2s Polyclonal Antibody

Catalog: HW027 100µl Orders: order@signalwayantibody.com







Application	Species Reactivity	Source	Molecular Wt.
WB IHC IF	Hu Mu Rt Other(Wide range)	Rabbit Polyconal Ab	11kDa

Description: Antibodies were produced by immunizing rabbits and

were purified by antigen affinity-chromatography.

Immunogen: A synthetic peptide corresponding to the amino terminus

of histone H4 in which Arg3 is di-methylated.

Formulation: Buffer: PBS with 0.02% sodium azide, 50% glycerol,

pH7.3.

Synonyms: H4; H4/n; H4F2; H4FN; FO108; HIST2H4

Accession No.: Gene ID: 8290 Swiss Prot: Q16695


Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in a histone cluster on chromosome 1. This gene is one of four histone genes in the cluster that are duplicated; this record represents the centromeric copy.


Application:

Recommended Dilutions:


WB 1:500 - 1:2000 IHC 1:50 - 1:200 IF 1:50 - 1:200



Western blot analysis of extracts of HeLa cell line and H4 protein expressed in E.coli., using H4R3me2s antibody.



Dot-blot analysis of all sorts of methylation peptidesusing H4R3me2s antibody.



Immunofluorescence analysis of 293T cell using H4R3me2s antibody. Blue: DAPI for nuclear staining.

This product is for in vitro research use only and is not intended for use in humans or animals.