**GRM8** Antibody

Catalog No: #36914

Description



Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Product Name	GRM8 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of total GRM8 protein.
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide corresponding to a region derived from internal residues of human glutamate receptor,
	metabotropic 8
Target Name	GRM8
Other Names	GLUR8; mGlu8; GPRC1H; MGLUR8
Accession No.	Swiss-Prot#: 000222NCBI Gene ID: 2918Gene Accssion: NP_000836
Uniprot	O00222
GeneID	2918;
SDS-PAGE MW	56kd
Concentration	2.4mg/ml
Formulation	Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
Storage	Store at -20°C

## Application Details

Western blotting: 1:500-1:2000

Immunohistochemistry: 1:50-1:200

## Images



Gel: 8%SDS-PAGE Lysates (from left to right): Mouse brain tissue Amount of lysate: 40ug per lane Primary antibody: 1/1200 dilution Secondary antibody dilution: 1/8000 Exposure time: 5 minutes



Immunohistochemical analysis of paraffin-embedded Human ovarain cancer tissue using #36914 at dilution 1/120.

## Background

L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.?

Note: This product is for in vitro research use only