CKMT1A/CKMT1B Antibody

Catalog No: #36996




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech@signalwayantibody.com
Product Name	CKMT1A/CKMT1B Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification.
Applications	WB IHC
Species Reactivity	Human, Mouse
Specificity	The antibody detects endogenous levels of total CKMT1A/CKMT1B protein.
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide of human CKMT1A/CKMT1B
Target Name	CKMT1A-CKMT1B
Other Names	CKMT; CKMT1; UMTCK
Accession No.	Swiss-Prot#: P12532NCBI Gene ID: 1159Gene Accssion: NP_066270
Uniprot	P12532
GeneID	1159;548596;
SDS-PAGE MW	47 kDa
Concentration	1 mg/ml
Formulation	Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
Storage	Store at -20°C

Application Details

Western blotting: 1:500-1:2000 Immunohistochemistry: 1:40-1:200

Images



Gel: 8% SDS-PAGE

Lysate: 40 ug


Lane: Mouse heart tissue lysate Primary antibody: (CKMT1A/CKMT1B

Antibody) at dilution 1/400

Secondary antibody: Goat anti rabbit IgG at 1/5000

dilution

Exposure time: 20 seconds



The image on the left is immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using (CKMT1A/CKMT1B Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: 200)

Background

Mitochondrial creatine (MtCK) kinase is responsible for the transfer of high energy phosphate from mitochondria to the cytosolic carrier, creatine. It belongs to the creatine kinase isoenzyme family. It exists as two isoenzymes, sarcomeric MtCK and ubiquitous MtCK, encoded by separate genes. Mitochondrial creatine kinase occurs in two different oligomeric forms: dimers and octamers, in contrast to the exclusively dimeric cytosolic creatine kinase isoenzymes. Many malignant cancers with poor prognosis have shown overexpression of ubiquitous mitochondrial creatine kinase; this may be related to high energy turnover and failure to eliminate cancer cells via apoptosis. Ubiquitous mitochondrial creatine kinase has 80% homology with the coding exons of sarcomeric mitochondrial creatine kinase. Two genes located near each other on chromosome 15 have been identified which encode identical mitochondrial creatine kinase proteins.?

Note: This product is for in vitro research use only