

ORC2 antibody

Catalog No: #22192

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	ORC2 antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC
Species Reactivity	Hu
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 208 and 449 of ORC2
Target Name	ORC2
Accession No.	Swiss-Prot:Q13416Gene ID:4999
Uniprot	Q13416
GeneID	4999;
Concentration	0.5mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 66kd

Western blotting: 1:500-1:3000

Immunohistochemistry: 1:100-1:250

Images




Sample (30 ug of whole cell lysate)

A: Raji

7.5% SDS PAGE

ORC2 antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded CL1-0 Xenograft, using ORC2 antibody at 1: 100 dilution.

Background

The origin recognition complex (ORC) is a highly conserved six subunits protein complex essential for the initiation of the DNA replication in eukaryotic cells. Studies in yeast demonstrated that ORC binds specifically to origins of replication and serves as a platform for the assembly of additional initiation factors such as Cdc6 and Mcm proteins. The protein encoded by this gene is a subunit of the ORC complex. This protein forms a core complex with ORC3L, -4L, and -5L. It also interacts with CDC45L and MCM10, which are proteins known to be important for the initiation of DNA replication. This protein has been demonstrated to specifically associate with the origin of replication of Epstein-Barr virus in human cells, and is thought to be required for DNA replication from viral origin of replication. [provided by RefSeq]

Note: This product is for in vitro research use only