IL-1F9 antibody

Catalog No: #22271

Description



## Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

| Description           |  |
|-----------------------|--|
| Product Name          | IL-1F9 antibody  |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | Purified by antigen-affinity chromatography.   |
| Applications          | WB IHC   |
| Species Reactivity    | Hu   |
| Immunogen Type        | Peptide  |
| Immunogen Description | Synthetic peptide contain a sequence corresponding to a region within amino acids 105 and 169 of Human |
|                       | IL1F9  |
| Target Name           | IL-1F9   |
| Accession No.         | Swiss-Prot:Q9NZH8Gene ID:56300   |
| Uniprot               | Q9NZH8   |
| GeneID                | 56300;   |
| Concentration         | 1mg/ml   |
| Formulation           | Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a       |
|                       | preservative.  |
| Storage               | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.              |
|                       |  |

## Application Details

| Predicted MW: 19kd               |  |
|----------------------------------|--|
| Western blotting: 1:500-1:3000   |  |
| Immunohistochemistry: 1:50-1:500 |  |

## Images



Sample(30 ug of whole cell lysate) A: H1299 B: HeLa S3 12% SDS PAGE Primary antibody diluted at 1: 500



Immunohistochemical analysis of paraffin-embedded hela xenograft, using IL1F9 antibody antibody at 1: 500 dilution.

## Background

The protein encoded by this gene is a member of the interleukin 1 cytokine family. The activity of this cytokine is mediated by interleukin 1 receptor-like 2 (IL1RL2/IL1R-rp2), and is specifically inhibited by interleukin 1 family, member 5 (IL1F5/IL-1 delta). Interferon-gamma, tumor necrosis factor-alpha and interleukin 1, beta (IL1B) are reported to stimulate the expression of this cytokine in keratinocytes. The expression of this cytokine in keratinocytes can also be induced by a contact hypersensitivity reaction or herpes simplex virus infection. This gene and eight other interleukin 1 family genes form a cytokine gene cluster on chromosome 2. [provided by RefSeq]

Note: This product is for in vitro research use only