

TAK1 (Phospho-Thr184) Antibody

Catalog No: #12153



Package Size: #12153-1 50ul #12153-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	TAK1 (Phospho-Thr184) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of TAK1 only when phosphorylated at threonine 184.
Immunogen Type	peptide
Immunogen Description	Peptide sequence around phosphorylation site of threonine 184 (I-Q-T(p)-H-M) derived from Human TAK1.
Target Name	TAK1
Modification	Phospho
Other Names	EC 2.7.11.25; M3K7; MAP3K7; Mitogen-activated protein kinase kinase kinase 7; TGF-beta- activated kinase 1; Transforming growth factor-beta-activated kinase 1; kinase TAK1
Accession No.	Swiss-Prot#:O43318;NCBI Gene#:6885
Uniprot	O43318
GeneID	6885;
SDS-PAGE MW	65kd
Concentration	1.0mg/ml
Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C


Application Details

Western blotting: 1:500~1:3000

Images



Western blot analysis of extracts from HepG2 cells, treated with TNF (20ng/ml, 5mins), using TAK1 (Phospho-Thr184) antibody #12153. The lane on the right is treated with the synthesized peptide.



Western blot analysis of extracts from 293 cells (Lane 2), using TAK1 (Phospho-Thr184) Antibody #12153. The lane on the left is treated with synthesized peptide.

Background

Serine/threonine kinase which acts as an essential component of the MAP kinase signal transduction pathway. Plays an important role in the cascades of cellular responses evoked by changes in the environment. Mediates signal transduction of TRAF6, various cytokines including interleukin-1 (IL-1), transforming growth factor-beta (TGFB), TGFB-related factors like BMP2 and BMP4, toll-like receptors (TLR), tumor necrosis factor receptor CD40 and B-cell receptor (BCR). Ceramides are also able to activate MAP3K7/TAK1. Once activated, acts as an upstream activator of the MKK/JNK signal transduction cascade and the p38 MAPK signal transduction cascade through the phosphorylation and activation of several MAP kinase kinases like MAP2K1/MEK1, MAP2K3/MKK3, MAP2K6/MKK6 and MAP2K7/MKK7.

Note: This product is for in vitro research use only