

## IGF-1R(Phospho-Tyr1165/Tyr1166) Antibody

Catalog No: #11088

Package Size: #11088-1 50ul #11088-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	IGF-1R(Phospho-Tyr1165/Tyr1166) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB, IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of IGF-1R only when phosphorylated at tyrosine 1165/tyrosine1166.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 1165/tyrosine 1166 (T-D-Y(p)-Y(p)-R-K) derived from Human IGF-1R .
Target Name	IGF-1R
Modification	Phospho
Other Names	Insulin-like growth factor I receptor; CD221; IGF1R; kinase IGF1R;
Accession No.	Swiss-Prot: P08069NCBI Protein: NP_000866.1
Uniprot	P08069
GeneID	3480;
Concentration	1.0mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


## Application Details

WB 1:500 - 1:1000, IHC 1 $\mu$ O 50 - 1 $\mu$ O 100

## Images



Western blot analysis of IGF1 Receptor(Phospho-Y1165/Y1166) expression in MCF7 (A),MCF7-insulin-15min (B), A2780 (C) whole cell lysates.(Predicted band size: 154 kD; Observed band size: 95; 200 kD)



Immunohistochemical analysis of IGF1 Receptor(Phospho-Y1165/Y1166) staining in human colorectal cancer formalin fixed paraffin embedded tissue section.

## Background

This receptor binds insulin-like growth factor 1 (IGF1) with a high affinity and IGF2 with a lower affinity. It has a tyrosine-protein kinase activity, which is necessary for the activation of the IGF1-stimulated downstream signaling cascade. When present in a hybrid receptor with INSR, binds IGF1.

Li S, et al. (1994) J Biol Chem; 269(51): 32558-64.

Note: This product is for in vitro research use only