AGT antibody

Catalog No: #38789

SAB Signalway Antibody

Package Size: #38789-1 50ul #38789-2 100ul Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


_				
	20	Cri	nti	ion
יע	-0	UH	νι	ווטו

Rabbit		
Polyclonal		
Antibodies were purified by affinity purification using immunogen.		
WB,IHC,IF		
Human,Mouse,Rat		
The antibody detects endogenous level of total AGT protein.		
Recombinant Protein		
Recombinant protein of human AGT.		
AGT		
ANHU; SERPINA8;		
Swiss-Prot#: P01019NCBI Gene ID: 183		
P01019		
183;		
53kd		
1.0mg/ml		
Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%		
sodium azide and 50% glycerol.		
Store at -20°C		


Application Details

WB□1:500 - 1:2000IHC□1:50 - 1:100IF□1:50 - 1:100


Images



Immunohistochemistry of paraffin-embedded human liver cancer using AGT at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded human placenta using AGT at dilution of 1:100 (40x lens).



Western blot analysis of extracts of HepG2 cells, using AGT at 1:1000 dilution.

Background

The protein encoded by this gene, pre-angiotensinogen or angiotensinogen precursor, is expressed in the liver and is cleaved by the enzyme renin in response to lowered blood pressure. The resulting product, angiotensin I, is then cleaved by angiotensin converting enzyme (ACE) to generate the physiologically active enzyme angiotensin II. The protein is involved in maintaining blood pressure and in the pathogenesis of essential hypertension and preeclampsia. Mutations in this gene are associated with susceptibility to essential hypertension, and can cause renal tubular dysgenesis, a severe disorder of renal tubular development. Defects in this gene have also been associated with non-familial structural atrial fibrillation, and inflammatory bowel disease.

Note: This product is for in vitro research use only