Product Datasheet

p53(Phospho-Ser315) Antibody

Catalog No: #11100

Description

Package Size: #11100-1 50ul #11100-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	p53(Phospho-Ser315) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC IF
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of p53 only when
	phosphorylated at serine 315.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 315 (S-S-S(p)-P-Q) derived from Human p53.
Target Name	p53
Modification	Phospho
Other Names	Tumor suppressor p53; Phosphoprotein p53; Antigen NY-CO-13; TP53;

Swiss-Prot: P04637NCBI Protein: NP_000537.3

sodium azide and 50% glycerol.

P04637 7157;

1.0mg/ml

Application Details

Accession No.

Uniprot

GeneID Concentration

Storage

Formulation

Predicted MW: 53kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%


Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.



Western blot analysis of extracts from Hela cells using p53(Phospho-Ser315) Antibody #11100(Lane 2) and the same antibody preincubated with blocking peptide(Lane1).



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using p53(Phospho-Ser315) Antibody #11100(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using p53(Phospho-Ser315) Antibody #11100.

Background

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over.

Lu, H. et al. (1997) Mol. Cell. Biol. 17, 5923-5934.

Lohrum, M. et.al. (1996) Oncogene 13, 2527-2539.

Posp

Note: This product is for in vitro research use only