SAP130 antibody

Catalog No: #22516

Description




Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Product Name	SAP130 antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC
Species Reactivity	Ни
Immunogen Type	Peptide
Immunogen Description	Synthetic peptide contain a sequence corresponding to a region within amino acids 1154 and 1217 of
	SAP130
Target Name	SAP130
Accession No.	Swiss-Prot:Q15393Gene ID:23450
Uniprot	Q15393
GeneID	23450;
Concentration	1mg/ml
Formulation	Supplied in 1XPBS, 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

## Application Details Predicted MW: 136kd Western blotting: 1:500-1:3000 Immunohistochemistry: 1:100-1:250

## Images



Sample (30 ug of whole cell lysate) A: Hela 5% SDS PAGE Primary antibody diluted at 1: 3000



Immunohistochemical analysis of paraffin-embedded Saos-2 xenograft, using SF3B3 antibody at 1: 100 dilution.

## Background

This gene encodes subunit 3 of the splicing factor 3b protein complex. Splicing factor 3b, together with splicing factor 3a and a 12S RNA unit, forms the U2 small nuclear ribonucleoproteins complex (U2 snRNP). The splicing factor 3b/3a complex binds pre-mRNA upstream of the intron's branch site in a sequence independent manner and may anchor the U2 snRNP to the pre-mRNA. Splicing factor 3b is also a component of the minor U12-type spliceosome. Subunit 3 has also been identified as a component of the STAGA (SPT3-TAF(II)31-GCN5L acetylase) transcription coactivator-HAT (histone acetyltransferase) complex, and the TFTC (TATA-binding-protein-free TAF(II)-containing complex). These complexes may function in chromatin modification, transcription, splicing, and DNA repair. [provided by RefSeq]

Note: This product is for in vitro research use only