## Tau(Phospho-Thr205) Antibody

Catalog No: #11108

Package Size: #11108-1 50ul #11108-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Tau(Phospho-Thr205) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of Tau only when phosphorylated at threonine 205.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 205 (P-G-T(p)-P-G) derived from Human Tau.
Target Name	Tau
Modification	Phospho
Other Names	MAPT; MTAPT; MTBT1; Neurofibrillary tangle protein; PHF-tau
Accession No.	Swiss-Prot: P10636NCBI Protein: NP _001116538.1
Uniprot	P10636
GeneID	4137;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.


Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

## **Application Details**


Predicted MW: 48 62 78 kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

## **Images**

Storage



Western blot analysis of extracts from mouse brain tissue using Tau(Phospho-Thr205) Antibody #11108 and the same antibody preincubated with blocking peptide.



Immunohistochemical analysis of paraffin-embedded rat hippocampal region tissue from a model with Alzheimer

## Background

Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.

del C Alonso A, et al. (2004) J Biol Chem; 279(33): 34873-81 Li G, Yin H, Kuret J (2004) J Biol Chem; 279(16): 15938-45 Giasson BI, et al. (2002) Biochemistry; 41(51): 15376-87

Liu F, et al. (2002) FEBS Lett; 530(1-3): 209

Taniguchi T, et al. (2001) J Biol Chem; 276(13): 10025-31

Note: This product is for in vitro research use only