Serine/threonine-protein kinase ATR

Catalog No: #AP79088



Package Size: #AP79088-1 50ug #AP79088-2 100ug #AP79088-3 1mg

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| D | e) | S | cri | p | tio | OI | n |
|---|----|---|-----|---|-----|----|---|
| | | | | | | | |

| Product Name | Serine/threonine-protein kinase ATR | | | |
|-----------------------|--------------------------------------------------------------------------------------------------------|--|--|--|
| Brief Description | Recombinant Protein | | | |
| Host Species | E.coli | | | |
| Purification | Greater than 90% by SDS-PAGE | | | |
| Species Reactivity | Mouse | | | |
| Immunogen Description | 2313-2558AA | | | |
| Other Names | Kiaa4069,Ataxia telangiectasia and Rad3-related protein | | | |
| Accession No. | Q9JKK8Gene name:Atr | | | |
| Uniprot | Q9JKK8 | | | |
| Calculated MW | 27.06 | | | |
| Tag Info | His | | | |
| Formulation | 50mM NaH2PO4, 500mM NaCl Buffer with 500mM Imidazole,10%glycerol(PH8.0) | | | |
| Storage | Store at -20C. (Avoid repeated freezing and thawing.)Repeated freezing and thawing is not recommended. | | | |
| | Store working aliquots at 4°C for up to one week. | | | |

Background

Serine/threonine protein kinase which activates checkpoint signaling upon genotoxic stresses such as ionizing radiation (IR), ultraviolet light (UV), or DNA replication stalling, thereby acting as a DNA damage sensor. Recognizes the substrate consensus sequence [ST]-Q. Phosphorylates BRCA1, CHEK1, MCM2, RAD17, RPA2, SMC1 and p53/TP53, which collectively inhibit DNA replication and mitosis and promote DNA repair, recombination and apoptosis. Phosphorylates Ser-139 of histone variant H2AX/H2AFX at sites of DNA damage, thereby regulating DNA damage response mechanism. Required for FANCD2 ubiquitination. Critical for maintenance of fragile site stability and efficient regulation of centrosome duplication (By similarity). Essential for preventing the occurrence of DNA damage during early embryogenesis.

References

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Note: This product is for in vitro research use only