

EPHA2 (phospho-Tyr588/596) Antibody

Catalog No: #13312

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	EPHA2 (phospho-Tyr588/596) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was purified by immunogen affinity chromatography.
Applications	WB IF
Species Reactivity	Hu,Ms,Rt
Specificity	Recognizes endogenous levels of EPHA2 (phospho-Tyr588/596) protein.
Immunogen Description	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human EPHA2.
Target Name	EPHA2; EPHA3; EPHA4
Other Names	EPHA2; ECK; Ephrin type-A receptor 2; Epithelial cell kinase; Tyrosine-protein kinase receptor ECK; EPHA3; ETK; ETK1; HEK; TYRO4; Ephrin type-A receptor 3; EPH-like kinase 4; EK4; hEK4; HEK; Human embryo kinase; Tyrosine-protein kinase TYRO4; Tyrosine-protein kinase receptor ETK1; Eph-like tyrosine kinase 1; EPHA4; HEK8; SEK; TYRO1; Ephrin type-A receptor 4; EPH-like kinase 8; EK8; hEK8; Tyrosine-protein kinase TYRO1; Tyrosine-protein kinase receptor SEK
Accession No.	Swiss-Prot#:P29317; P29320; P54764NCBI Gene ID:1969; 2042; 2043
Uniprot	P29317
GeneID	1969;
Calculated MW	130KD
Concentration	1 mg/ml
Formulation	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C


Application Details

Western blotting:1:500 - 1:1000Immunofluorescence:1:100 - 1:300

Images



Western blot analysis of EPHA2 (phospho-Tyr588/596) expression in A431, mouse brain whole cell lysates.



Immunofluorescent analysis of EPHA2 (phospho-Tyr588/596) staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Note: This product is for in vitro research use only