CLIP3 Antibody

Catalog No: #47818




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	Support: tech@signalwayantibody.com
Product Name	CLIP3 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was purified by immunogen affinity chromatography.
Applications	WB, IHC, IF/ICC
Species Reactivity	Hu,Ms,Rt
Specificity	Recognizes endogenous levels of CLIP3 protein.
Immunogen Description	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CLIP3.
Target Name	CLIP3
Other Names	CLIPR59; CAP-Gly domain-containing linker protein 3; Cytoplasmic linker protein 170-related 59 kDa protein;
	CLIP-170-related 59 kDa protein; CLIPR-59
Accession No.	Swiss-Prot#:Q96DZ5NCBI Gene ID:25999
Uniprot	Q96DZ5
GeneID	25999;
Calculated MW	60KD
Concentration	1 mg/ml
Formulation	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium
	azide.
Storage	Store at -20°C


Application Details

WB (1/500 - 1/1000), IHC (1/100 - 1/200), IF/ICC (1/100 - 1/500)


Images



Western blot analysis of CLIP3 expression in human brain (A), A549 (B) whole cell lysates



Immunohistochemical analysis of CLIP3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CLIP3 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Note: This product is for in vitro research use only