NFYB Antibody

Catalog No: #47839




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name NFYB Antibody Host Species Rabbit Clonality Polyclonal Purification The antibody was purified by immunogen affinity chromatography. Applications WB IHC Species Reactivity Hu,Ms,Rt Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Storage Store at -20°C	Description	Support: tech@signalwayantibody.com
Clonality Polyclonal Purification The antibody was purified by immunogen affinity chromatography. Applications WB IHC Species Reactivity Hu,Ms,Rt Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Product Name	NFYB Antibody
Purification The antibody was purified by immunogen affinity chromatography. Applications WB IHC Species Reactivity Hu,Ms,Rt Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCI, 0.02% sodium azide and 50% glycerol.	Host Species	Rabbit
Applications WB IHC Species Reactivity Hu,Ms,Rt Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Clonality	Polyclonal
Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Purification	The antibody was purified by immunogen affinity chromatography.
Specificity Recognizes endogenous levels of NFYB protein. Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Applications	WB IHC
Immunogen Description KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB. Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Species Reactivity	Hu,Ms,Rt
Target Name NFYB Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Specificity	Recognizes endogenous levels of NFYB protein.
Other Names HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Immunogen Description	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NFYB.
transcription factor Y subunit B; NF-YB Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Target Name	NFYB
Accession No. Swiss-Prot#:P25208NCBI Gene ID:4801 Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Other Names	HAP3; Nuclear transcription factor Y subunit beta; CAAT box DNA-binding protein subunit B; Nuclear
Uniprot P25208 GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.		transcription factor Y subunit B; NF-YB
GeneID 4801; Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Accession No.	Swiss-Prot#:P25208NCBI Gene ID:4801
Calculated MW 32KD Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Uniprot	P25208
Concentration 1 mg/ml Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	GeneID	4801;
Formulation Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.	Calculated MW	32KD
and 50% glycerol.	Concentration	1 mg/ml
	Formulation	Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide
Storage Store at -20°C		and 50% glycerol.
	Storage	Store at -20°C


Application Details

Western blotting:1:500 - 1:1000lmmunohistochemistry:1:50 - 1:100

Images



Western blot analysis of MRPL22 expression in Hela, HepG2 whole cell lysates.



Immunohistochemical analysis of NFYB staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note: This product is for in vitro research use only