Maltose Binding Protein Rabbit mAb

Catalog No: #48666

Package Size: #48666-1 50ul #48666-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | Maltose Binding Protein Rabbit mAb |
|-----------------------|---|
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal antibody |
| Clone No. | SR41-04 |
| Purification | ProA affinity purified |
| Applications | WB |
| Species Reactivity | Escherichia coli |
| Immunogen Description | recombinant protein |
| Other Names | ECK4026 antibody JW3994 antibody Mal E antibody MalE antibody malJ antibody Maltodextrin binding |
| | protein antibody Maltose ABC transporter periplasmic protein antibody Maltose binding periplasmic protein |
| | antibody MMBP antibody Periplasmic maltose binding protein antibody |
| Accession No. | Swiss-Prot:P0AEX9Gene ID:948538 |
| Uniprot | P0AEX9 |
| GenelD | 948538; |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |
| | |

Application Details

WB: 1:1,000-5,000

Images



Western blot analysis of Maltose Binding Protein on recombinant MBP-tag protein lysates using anti-Maltose Binding Protein antibody at 1/1,000 dilution.

Background

Plasmid vectors for the expression of coding regions of eukaryotic genes in bacterial, insect and mammalian hosts are in common usage; such expression vectors frequently encode hybrid fusion proteins consisting in part of prokaryotic and in part, eukaryotic specified proteins. One such system utilizes maltose binding protein (MBP), the 370 amino acid product of the E. coli mal E gene. Plasmid vectors have been constructed utilizing

the MBP domain that allow the synthesis of high levels of MBP-fusion proteins that can be purified in a one step procedure by affinity chromatography cross linked amylose resin. Once bound to amylose, the MBP protein can then be separated from the target protein by cleavage by coagulation factor Xa at a specific four residue site. Alternatively, the intact fusion protein can be specifically eluted from the resin by the addition of excess free maltose. Subsequent to elution, MBP fusion protein can be visualized either by Western blot analysis or immunoprecipitation using antibodies specific for the MBP-tag. Expression systems utilizing the MBP fusion tag include pCG-806fx and pMal vectors.

References

Note: This product is for in vitro research use only