Caspase-1 Rabbit mAb

Catalog No: #48847

Package Size: #48847-1 50ul #48847-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$				
	00	cri	ntı	on
	-	CIL	UII	ULL
	\sim		~	

Product Name	Caspase-1 Rabbit mAb			
Host Species	Recombinant Rabbit			
Clonality	Monoclonal antibody			
Clone No.	SU40-07			
Purification	ProA affinity purified			
Applications	WB, IHC, FC			
Species Reactivity	Hu, Ms, Rt			
Immunogen Description	recombinant protein			
Other Names	CASP-1 antibody CASP1 antibody CASP1_HUMAN antibody Caspase 1 antibody Caspase-1 subunit p10			
	antibody ICE antibody IL-1 beta-converting enzyme antibody IL-1BC antibody IL1 beta converting enzyme			
	antibody IL1B convertase antibody Interleukin 1 beta convertase antibody Interleukin 1B converting enzyme			
	antibody Interleukin-1 beta convertase antibody Interleukin-1 beta-converting enzyme antibody p45 antibody			
Accession No.	Swiss-Prot#:P29466			
Uniprot	P29466			
GeneID	834;			
Calculated MW	45 kDa			
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.			
Storage	Store at -20°C			


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of Caspase-1 on different lysates using anti-Caspase-1 antibody at 1/1,000 dilution. Positive control: Lane 1: THP-1 Lane 2: Jurkat




Immunohistochemical analysis of paraffin-embedded human lung tissue using anti-Caspase-1 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue using anti-Caspase-1 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue using anti-Caspase-1 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of Hela cells with Caspase-1 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Caspase-1, originally designated ICE (for IL-1 converting enzyme), is a member of the group of caspases with large prodomains. Caspase-1 promotes maturation of interleukin IL-1 β and interleukin18 (IL-18) by proteolytic cleavage of precursor forms into biologically active pro-inflamatory cytokines. Active caspase-1, a (p20/p10)2 tetramer, is necessary and sufficient for cleavage of precursor IL-1 as well as for induction of apoptosis in some cell lines. The highly conserved family of caspases mediate many of the morphological and biochemical features of apoptosis, including structural dismantling of cell bodies and nuclei, fragmentation of genomic DNA, destruction of regulatory proteins and propagation of other pro-apoptotic molecules. The human Caspase-1 gene maps to chromosome 2q14 and encodes a cytoplasmic protein expressed in liver, heart, skeletal muscle kidney and testis. Caspase-1 has been implicated in inflammation, septic shock, and other situations such as wound healing and the growth of certain

leuker	nias.			

References

Note: This product is for in vitro research use only