

Phospho-STAT1(S727) Rabbit mAb

Catalog No: #13392

Package Size: #13392-1 50ul #13392-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	Phospho-STAT1(S727) Rabbit mAb
Clone No.	SN67-04
Purification	ProA affinity purified
Applications	WB, IHC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser727 of human STAT1.
Other Names	Signal transducer and activator of transcription 1 91kD antibody CANDF7 antibody DKFZp686B04100 antibody IMD31A antibody IMD31B antibody IMD31C antibody ISGF 3 antibody ISGF-3 antibody OTTHUMP00000163552 antibody OTTHUMP00000165046 antibody OTTHUMP00000165047 antibody OTTHUMP00000205845 antibody Signal transducer and activator of transcription 1 91kDa antibody Signal transducer and activator of transcription 1 antibody Signal transducer and activator of transcription 1, 91kD antibody Signal transducer and activator of transcription 1-alpha/beta antibody STAT 1 antibody Stat1 antibody STAT1_HUMAN antibody STAT91 antibody Transcription factor ISGF 3 components p91 p84 antibody Transcription factor ISGF-3 components p91/p84 antibody
Accession No.	Swiss-Prot#:P42224
Uniprot	P42224
GenelD	6772;
Calculated MW	87 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details


WB: 1:1,000

IHC: 1:50-1:200


Images



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-Phospho-STAT1(S727) antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-Phospho-STAT1(S727) antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse colon tissue using anti-Phospho-STAT1(S727) antibody. Counter stained with hematoxylin.

Background

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of Jak kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

References

Note: This product is for in vitro research use only