

Tyrosine Hydroxylase Rabbit mAb

Catalog No: #49178



Package Size: #49178-1 50ul #49178-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Tyrosine Hydroxylase Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SD080-02
Purification	ProA affinity purified
Applications	WB, ICC, IHC, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Other Names	Dystonia 14 antibody DYT14 antibody DYT5b antibody EC 1.14.16.2 antibody OTTHUMP00000011225 antibody OTTHUMP00000011226 antibody ple antibody Protein Pale antibody TH antibody The antibody TY3H_HUMAN antibody TYH antibody Tyrosine 3 hydroxylase antibody Tyrosine 3 monooxygenase antibody Tyrosine 3-hydroxylase antibody Tyrosine 3-monooxygenase antibody Tyrosine hydroxylase antibody
Accession No.	Swiss-Prot#:P07101
Uniprot	P07101
GeneID	7054;
Calculated MW	59 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:50,000-1:100,000IHC: 1:50-1:200 ICC: 1:100-1:500FC: 1:50-1:100


Images




Western blot analysis of Tyrosine Hydroxylase on different lysates using anti-Tyrosine Hydroxylase antibody at 1/50,000 dilution. Positive control: Lane 1: PC-12 Lane 2: Mouse brain




Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti-Tyrosine Hydroxylase antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-Tyrosine Hydroxylase antibody. Counter stained with hematoxylin.



ICC staining Tyrosine Hydroxylase in N2A cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Tyrosine Hydroxylase in SH-SY-5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of SH-SY-5Y cells with Tyrosine Hydroxylase antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody

Background

The enzyme tyrosine hydroxylase (TH), also designated tyrosine 3-monooxygenase (TY3H), catalyzes the conversion of tyrosine to L-dopa, which is the rate limiting step in the biosynthesis of catecholamines such as dopamine, adrenalin and noradrenalin. TH is thought to play a role in the pathogenesis of Parkinsons disease, which is associated with reduced dopamine levels. Two transcription factor binding sites in the proximal region of the TH gene, the TPA-responsive element (TRE) and the c-AMP responsive element (CRE), have been implicated in the complex regulation of the TH gene. TH is also known to be upregulated by the glia maturation factor (GMF), a Cdc 10/SWI6 motif-containing protein called V-1, and a variety of additional compounds.

References

Note: This product is for in vitro research use only