

HDAC10 Rabbit mAb

Catalog No: #49182




Package Size: #49182-1 50ul #49182-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description	
Product Name	HDAC10 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SD08-11
Purification	ProA affinity purified
Applications	WB, ICC, IP
Species Reactivity	Hu, Ms
Immunogen Description	recombinant protein
Other Names	DKFZP761B039 antibody HD 10 antibody HD10 antibody HDA10_HUMAN antibody HDAC 10 antibody Hdac10 antibody Histone deacetylase 10 antibody MGC149722 antibody OTTHUMP00000028555 antibody
Accession No.	Swiss-Prot#:Q969S8
Uniprot	Q969S8
GeneID	83933;
Calculated MW	43 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details
WB: 1:1,000ICC: 1:50-1:200


Images




Western blot analysis of HDAC10 on HepG2 cells lysates using anti-HDAC10 antibody at 1/1,000 dilution.



ICC staining HDAC10 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining HDAC10 in A431 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining HDAC10 in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Histone deacetylases (HDACs) play an important role in the modification of chromatin structure and thus in the suppression and activation of transcription and cellular differentiation. There are 11 members in the HDAC family that are divided into four classes. Class I HDACs represent homologs of the yeast histone deacetylase Rpd3, class II HDACs share strong homology with the yeast histone deacetylase Hda1, class III HDACs are closely related to the yeast Sir2 protein and class IV HDACs comprise histone deacetylase 11 (HDAC11)-related enzymes. HDAC10, also known as HD10, is a member of the class II HDACs. It contains an N-terminal Hda1p-related catalytic domain and a unique C-terminal leucine-rich domain. HDAC10 is ubiquitously expressed and can shuttle between the cytoplasm and nucleus in response to cellular signals. It is able to repress transcription and, like other class II HDAC members, its enzymatic activity is inhibited by Trichostatin A (TSA).

References

Note: This product is for in vitro research use only