

STAT4 Rabbit mAb

Catalog No: #49252



Package Size: #49252-1 50ul #49252-2 100ul


Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

Description	
Product Name	STAT4 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JJ08-85
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Other Names	Signal transducer and activator of transcription 4 antibody SLEB11 antibody STAT4 antibody STAT4_HUMAN antibody
Accession No.	Swiss-Prot#:Q14765
Uniprot	Q14765
GeneID	6775;
Calculated MW	86 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details	
WB: 1:1,000IHC: 1:50-1:200	ICC: 1:100-1:500FC: 1:50-1:100

Images


Western blot analysis of STAT4 on Daudi cells lysates using anti-STAT4 antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-STAT4 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human pancreas tissue using anti-STAT4 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue using anti-STAT4 antibody. Counter stained with hematoxylin.




ICC staining STAT4 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining STAT4 in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining STAT4 in SW480 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Hela cells with STAT4 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of Jak kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

References

Note: This product is for in vitro research use only