

## NQO1 Rabbit mAb

Catalog No: #49364

Package Size: #49364-1 50ul #49364-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	NQO1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JF440-1
Purification	ProA affinity purified
Applications	WB, ICC/IF, IP, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Other Names	Azoreductase antibody Cytochrome b 5 reductase antibody DHQU antibody DIA 4 antibody DIA4 antibody Diaphorase (NADH/NADPH) (cytochrome b 5 reductase) antibody Diaphorase (NADH/NADPH) antibody Diaphorase 4 antibody Dioxin inducible 1 antibody DT diaphorase antibody DT-diaphorase antibody DTD antibody Menadione reductase antibody NAD(P)H dehydrogenase [quinone] 1 antibody NAD(P)H dehydrogenase quinone 1 antibody NAD(P)H menadione oxidoreductase 1 dioxin inducible antibody NAD(P)H: menadione oxidoreductase 1 dioxin inducible 1 antibody NAD(P)H:menadione oxidoreductase 1 antibody NAD(P)H:Quinone acceptor oxidoreductase type 1 antibody NAD(P)H:quinone oxidoreductase 1 antibody NAD(P)H:quinone oxireductase antibody NMOR 1 antibody NMOR I antibody NMOR1 antibody NMORI antibody NQO 1 antibody NQO1 antibody NQO1_HUMAN antibody Phylloquinone reductase antibody QR 1 antibody QR1 antibody Quinone reductase 1 antibody
Accession No.	Swiss-Prot#:P15559
Uniprot	P15559
GeneID	1728;
Calculated MW	31/27 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:1,000-5,000 ICC: 1:50-1:200 IP: 1:50 FC: 1:50-1:100

## Images



Western blot analysis of NQO1 on different lysates using anti-NQO1 antibody at 1/1,000 dilution. Positive control:  
Lane 1: SH-SY-5Y    Lane 2: Mouse kidney    Lane 3: Raji



Flow cytometric analysis of HeLa cells with NQO1 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody

## Background

NAD(P)H:quinone oxidoreductase 1 (NQO1) and NRH:quinone oxidoreductase (NQO2) are flavoproteins that catalyze the metabolic detoxification of quinones and their derivatives to hydroquinones, using either NADH or NADPH as the electron donor. This protects cells against quinone-induced oxidative stress, cytotoxicity, and mutagenicity. Many tumors overexpress NQO1, which is an obligate two-electron reductase that deactivates toxins and activates bio-reductive anticancer drugs. NQO1, a 274 amino acid protein, is ubiquitously expressed, but the expression level varies among tissues. NQO1 gene expression is coordinately induced in response to xenobiotics, antioxidants, heavy metals and radiation. The antioxidant response element (ARE) in the NQO1 gene promoter is essential for expression and coordinated induction of NQO1. ARE activation by tert-butylhydroquinone is dependent on PI3-kinase, which lies upstream of Nrf2. Nrf2, c-Jun, Nrf1, Jun-B and Jun-D bind to the ARE and regulate expression and induction of NQO1 gene. Maf-Maf homodimers and possibly Maf-Nrf2 heterodimers play a role in negative regulation of ARE-mediated transcription, but Maf-Nrf1 heterodimers fail to bind with the NQO1 gene ARE and do not repress NQO1 transcription.

## References

Note: This product is for in vitro research use only