

Endo G Rabbit mAb

Catalog No: #49763



Package Size: #49763-1 50ul #49763-2 100ul

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)


Description

Product Name	Endo G Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JU31-30
Purification	ProA affinity purified
Applications	WB,ICC
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Other Names	EndoG antibody    Endonuclease G antibody    Endonuclease G mitochondrial antibody    EndonucleaseG antibody    FLJ27463 antibody    Mitochondrial endonuclease G antibody    NUCG_HUMAN antibody
Accession No.	Swiss-Prot#:Q14249
Uniprot	Q14249
GeneID	2021;
Calculated MW	36 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:1,000    ICC: 1:50-1:200


Images




Western blot analysis of Endo G on mouse heart tissue lysate using anti-Endo G antibody at 1/500 dilution.



ICC staining Endo G in PC-3M cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Endo G in SH-SY5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Endo G in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised

## Background

Endonuclease G (ENDOG), a nuclear encoded protein, localizes to the mitochondria. This sugar-nonspecific nuclease, responsible for major mitochondrial nuclease activity, preferentially cleaves single-stranded DNA(ssDNA). Synthesized as a propeptide with an amino-terminal presequence that targets the nuclease to mitochondria, ENDOG translocates to nuclei on apoptotic stimulation and act as a nuclease without sequence specificity. Both exonucleases and DNase I stimulate the ability of ENDOG to generate double-stranded DNA cleavage products at physiological ionic strengths, suggesting that these activities work in concert with ENDOG in apoptotic cells to ensure efficient DNA breakdown. In addition to deoxyribonuclease activities, ENDOG also has ribonuclease (RNase) and RNase H activities. ENDOG is capable of generating the RNA primers required by DNA polymerase gamma to initiate replication of mitochondrial DNA. ENDOG exists in the mitochondrial intermembrane space, but not in the matrix where mtDNA replication occurs. This enzyme provides an important nicking function for mitochondrial DNA specifically cleaving DNA at GC tracts. Human ENDOG maps to chromosome 9q34.11.

## References

Note: This product is for in vitro research use only