

Matrin 3 Rabbit mAb

Catalog No: #49779



Package Size: #49779-1 50ul #49779-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Matrin 3 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JU93-43
Purification	ProA affinity purified
Applications	WB,ICC,IF,IHC,FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Recombinant protein
Other Names	ALS21 antibody KIAA0723 antibody Matr3 antibody MATR3_HUMAN antibody Matrin-3 antibody Matrin3 antibody MPD2 antibody VCPDM antibody Vocal cord and pharyngeal weakness with distal myopathy antibody
Accession No.	Swiss-Prot#:P43243
Uniprot	P43243
GeneID	9782;
Calculated MW	95 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of Matrin 3 on different lysates using anti-Matrin 3 antibody at 1/1,000 dilution. Positive control:
Lane 1: Daudi Lane 2: Mouse brain tissue




Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti-Matrin 3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Matrin 3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue using anti-Matrin 3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human liver tissue using anti-Matrin 3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue using anti-Matrin 3 antibody. Counter stained with hematoxylin.




ICC staining Matrin 3 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Matrin 3 in LOVO cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Matrin 3 in SH-SY-5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of LOVO cells with Matrin 3 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

Background

Matrin-3 is a nuclear matrix protein containing one matrin-type zinc finger and two RRM (RNA recognition motif) domains. Matrin-3 plays a role in transcription and may interact with other nuclear matrix proteins to form the internal fibrogranular network. In association with the PSF-p54/NRB heteromer, Matrin-3 may play a role in the nuclear retention of defective RNAs. As the main substrate for PKA-mediated phosphorylation, Matrin-3 may serve as a rapid way of transferring information from synapses containing NMDA receptors to neuronal nuclei under physiological conditions. Also, the phosphorylation of Matrin-3 may contribute to neuronal death under pathological conditions. It is likely that Matrin-3 activity is regulated by calcium dependent interaction with CaM I and also by caspase induced cleavage.

References

Note: This product is for in vitro research use only