

Arp3 Rabbit mAb

Catalog No: #49873



Package Size: #49873-1 50ul #49873-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Arp3 Rabbit mAb			
Host Species	Recombinant Rabbit			
Clonality	Monoclonal antibody			
Clone No.	JB33-44			
Purification	ProA affinity purified			
Applications	WB,ICC,IF,IHC,FC			
Species Reactivity	Hu, Ms, Rt			
Immunogen Description	Recombinant protein of C-terminal human Arp3.			
Other Names	Actin like protein 3 antibody Actin related protein 3 antibody ACTR3 antibody ARP3 actin related protein 3 homolog (yeast) antibody			
Accession No.	Swiss-Prot#:P61158			
Uniprot	P61158			
GeneID	10096;			
Calculated MW	47 kDa			
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.			
Storage	Store at -20°C			


Application Details

WB: 1:500-1:1,000
IHC: 1:50-1:200
ICC: 1:50-1:200
FC: 1:50-1:100


Images




Western blot analysis of Arp3 on different lysates using anti-Arp3 antibody at 1/500 dilution. Positive control: Lane 1: A431 Lane 2: Mouse placenta Lane 3: Mouse thymus




Flow cytometric analysis of HL-60 cells with Arp3 antibody at 1/100 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.




Western blotting analysis using Arp3 Antibody #49873.




#49873 at 1/100 staining Human stomach cancer by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




#49873 at 1/100 staining rat kidney by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.




#49873 at 1/100 staining mouse kidney by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the primary antibody at 4°C overnight. An HRP conjugated anti-Rabbit antibody was used as the secondary antibody.



ICC staining Arp3 in HUVEC cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Arp3 in LOVO Cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Arp3 in SiHa Cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Actin polymerization is required for a variety of cell functions, including chemotaxis, cell migration, cell adhesion, and platelet activation. Cells trigger actin polymerization through either the de novo nucleation of filaments from monomeric actin, the severing of existing filaments to create uncapped barbed ends, or the uncapping existing barbed ends. The nucleation of actin is a rate-limiting and unfavorable reaction in actin polymerization and therefore requires the involvement of the Arp2/3 complex, which helps create new filaments and promotes the end-to-side cross-linking of actin filaments into the branching meshwork. The Arp2/3 complex consists of the actin-related proteins Arp2 and Arp3, and various other accessory proteins. The Arp2/3 complex promotes actin nucleation by binding the pointed end of actin filaments, or by associating with the side of an existing filament, and nucleates growth in the barbed direction. In addition, the Arp2/3 complex also mediates actin cytoskeletal outgrowths that are regulated by the Rho family of small GTPases. In response to GTP-binding Cdc42, the Arp2/3 complex binds the Cdc42 substrates, namely the WASP proteins, and initiates the formation of lamellipodia and filopodia.

Note: This product is for in vitro research use only