

Bovine Pyrroline-5-carboxylate reductase 1, mitochondrial (PYCR1) ELISA Kit

Catalog No: #EK7909

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #EK7909-1 48T #EK7909-2 96T

Description

Product Name	Bovine Pyrroline-5-carboxylate reductase 1, mitochondrial (PYCR1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Bovine (Bos taurus; Cattle)
Other Names	ARCL2B; P5C; P5CR; PIG45; PP222; PRO3; PYCR; P5C reductase proliferation-inducing protein 45
Accession No.	Q58DT4
Uniprot	Q58DT4
GeneID	539606;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate PYCR1 in samples. An antibody specific for PYCR1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyPYCR1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PYCR1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PYCR1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Product Overview:PYCR1 catalyzes the NAD(P)H-dependent conversion of pyrroline-5-carboxylate to proline. This enzyme may also play a physiologic role in the generation of NADP(+) in some cell types. The protein forms a homopolymer and localizes to the mitochondrion. Alternate splicing results in two transcript variants encoding different isoforms.The 1,810-bp cDNA hybridized to a 1.85-kb mRNA in samples from human cell lines and predicated a 319-amino acid, 33.4-kD protein.

PYCR1 protein colocalized with various mitochondrial markers, including delta-1-pyrroline-5-carboxylate synthetase (P5CS), which also belongs to the proline metabolic pathway. Knockdown of the orthologous genes in Xenopus and zebrafish led to epidermal hypoplasia and blistering that was accompanied by a massive increase in apoptosis.

Note: This product is for in vitro research use only