

# Mouse Next to BRCA1 gene 1 protein (NBR1) ELISA Kit

Catalog No: #EK8802

Package Size: #EK8802-1 48T #EK8802-2 96T

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)  
Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Mouse Next to BRCA1 gene 1 protein (NBR1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	1A1-3B; FLJ55359; FLJ98272; KIAA0049; M17S2; MIG19; membrane component; chromosome 17; surface marker 2 (ovarian carcinoma antigen CA125) migration-inducing protein 19
Accession No.	P97432
Uniprot	P97432
GeneID	17966;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:Request Information
Sensitivity:Request Information
Sample Type:Serum, Plasma, Other biological fluids
Sample Volume: 1-200 µL
Assay Time:1-4.5h
Detection wavelength:450 nm

## Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate NBR1 in samples. An antibody specific for NBR1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyNBR1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for NBR1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of NBR1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only