

## Dhh antibody

Catalog No: #22723

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description


|                       |   |
|-----------------------|---|
| Product Name          | Dhh antibody  |
| Host Species          | Rabbit  |
| Clonality             | Polyclonal  |
| Purification          | Purified by antigen-affinity chromatography.  |
| Applications          | WB  |
| Species Reactivity    | Hu  |
| Immunogen Type        | Recombinant protein   |
| Immunogen Description | Recombinant protein fragment contain a sequence corresponding to a region within amino acids 135 and 388 of Dhh |
| Target Name           | Dhh   |
| Accession No.         | Swiss-Prot:O43323Gene ID:50846  |
| Uniprot               | O43323  |
| GeneID                | 50846;  |
| Concentration         | 1mg/ml  |
| Formulation           | Supplied in 0.1M Tris-buffered saline with 20% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative.  |
| Storage               | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.                       |

## Application Details

Predicted MW: 44kd

Western blotting: 1:500-1:3000

## Images



Sample (30 ug of whole cell lysate)  
A: A431  
10% SDS PAGE  
Primary antibody diluted at 1: 10000

## Background

This gene encodes a member of the Hedgehog family. The hedgehog gene family encodes signaling molecules that play an important role in regulating morphogenesis. This protein is predicted to be made as a precursor that is autocatalytically cleaved; the N-terminal portion is soluble and

contains the signalling activity while the C-terminal portion is involved in precursor processing. More importantly, the C-terminal product covalently attaches a cholesterol moiety to the N-terminal product, restricting the N-terminal product to the cell surface and preventing it from freely diffusing throughout the organism. Defects in this protein have been associated with partial gonadal dysgenesis (PGD) accompanied by minifascicular polyneuropathy. This protein may be involved in both male gonadal differentiation and perineurial development. [provided by RefSeq]

---

Note: This product is for in vitro research use only