

Mouse DIS3-like exonuclease 2 (DIS3L2) ELISA Kit

Catalog No: #EK10669



Package Size: #EK10669-1 48T #EK10669-2 96T

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

| | |
|--------------------|--|
| Product Name | Mouse DIS3-like exonuclease 2 (DIS3L2) ELISA Kit |
| Brief Description | ELISA Kit |
| Applications | ELISA |
| Species Reactivity | Mouse (Mus musculus) |
| Other Names | FAM6A; FLJ36974; MGC42174; OTTHUMP00000203808 family with sequence similarity 6; member A |
| Accession No. | Q8CI75 |
| Uniprot | Q8CI75 |
| GeneID | 208718; |
| Storage | <p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p> |

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate DIS3L2 in samples. An antibody specific for DIS3L2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyDIS3L2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for DIS3L2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of DIS3L2 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Product Overview:The exosome is a multisubunit complex composed of several highly conserved subunits, some of which are 3' to 5' exoribonucleases. The complex is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich (ARE) elements in their untranslated 3' region.

DIS3L2 (DIS3-like exonuclease 2) is an 885 amino acid protein that is thought to function as an exonuclease and may be required for the 3' processing of pre-mRNA into mature mRNA. Defects or chromosomal translocations involving the gene encoding DIS3L2 may be associated with Marfanoid habitus, a genetic disorder characterized by abnormalities in the skeleton, eyes and cardiovascular system. DIS3L2 is expressed as five isoforms due to alternative splicing events.

Note: This product is for in vitro research use only