

Mouse Myosin-4 (MYH4) ELISA Kit

Catalog No: #EK11210



Package Size: #EK11210-1 48T #EK11210-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Mouse Myosin-4 (MYH4) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	MYH2B; MyHC-2B; MyHC-IIb; myosin; heavy polypeptide 4; skeletal muscle
Accession No.	Q5SX39
Uniprot	Q5SX39
GeneID	17884;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:15.6-1000 pg/mL

Sensitivity:6.4 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MYH4 in samples. An antibody specific for MYH4 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMYH4 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MYH4 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MYH4 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Product Overview:All MHC proteins share approximately 100% conservation in the phosphate-binding loop, several helices forming the nucleotide-binding pocket, residues involved in actin binding, residues involved in the stereospecific hydrophobic rigor state interaction of actin with myosin, a cleft that divides the ATP- and actin-binding sites, and a converter domain, or fulcrum, containing 2 conserved cysteines.the MYH4 gene would appear to be located on 17p13.1, the location of 4 other myosin heavy chain genes (MYH1; MYH2; MYH3; and MYH8). Furthermore, Soussi-Yanicostas et al. (1993) had evidence from analysis of YACs that the embryonic and fetal genes, on the one hand, and the adult fast myosin heavy chain genes, on the other hand, are contained within different genomic fragments.

Note: This product is for in vitro research use only