Rho guanine nucleotide exchange factor 7 Polyclonal Antibody

SAB Signalway Antibody

Catalog No: #42257

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | |
|------------------|-------|-----|----|
| | escri | nti | n |
| \boldsymbol{L} | COUL | ่บแ | UH |

| Product Name | Rho guanine nucleotide exchange factor 7 Polyclonal Antibody | |
|-----------------------|--|--|
| Host Species | Rabbit | |
| Clonality | Polyclonal | |
| Purification | Caprylic Acid Ammonium Sulfate Precipitation purified | |
| Applications | WB | |
| Species Reactivity | Hu | |
| Specificity | The antibody detects endogenous level of total Rho guanine nucleotide exchange factor 7 polyclonal antibody. | |
| Immunogen Type | protein | |
| Immunogen Description | unogen Description Recombinant Human Rho guanine nucleotide exchange factor 7 protein | |
| Target Name | Rho guanine nucleotide exchange factor 7 | |
| Other Names | Beta-Pix, COOL-1, PAK-interacting exchange factor beta, p85, ARHGEF7, COOL1, KIAA0142, P85SPR, | |
| | PAK3BP, PIXB ORF Names: Nbla10314 | |
| Accession No. | Swiss-Prot#: Q14155 | |
| Uniprot | Q14155 | |
| GeneID | 8874; | |
| Calculated MW | 90kd | |
| Formulation | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 | |
| Storage | Store at -20°C | |
| | | |

Application Details

Western blotting: □1:500 - 1:1000

Images



All lanes : Rho guanine nucleotide exchange factor 7 antibody

at 2ug/ml

Lane 1 : mouse skeletal muscle tissue Lane 2 : 293T whole cell lysate

Secondary

Goat polyclonal to Rabbit IgG at 1/10000 dilution

Predicted band size:90KDa Observed band size:90KDa

Background

Acts as a RAC1 guanine nucleotide exchange factor (GEF) and can induce membrane ruffling. Functions in cell migration, attachment and cell

spreading. Promotes targeting of RAC1 to focal adhesions By similarity. May function as a positive regulator of apoptosis. Downstream of NMDA receptors and CaMKK-CaMK1 signaling cascade, promotes the formation of spines and synapses in hippocampal neurons.

References

[1] Prediction of the coding sequences of unidentified human genes. IV. The coding sequences of 40 new genes (KIAA0121-KIAA0160) deduced by analysis of cDNA clones from human cell line KG-1.Nagase T., Seki N., Tanaka A., Ishikawa K., Nomura N.DNA Res. 2:1

Note: This product is for in vitro research use only