

## Human Amiodarone (AD) ELISA Kit

Catalog No: #EK12140



Package Size: #EK12140-1 48T #EK12140-2 96T

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Human Amiodarone (AD) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:31.25-2000 pg/mL

Sensitivity:13.7 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200  $\mu$ L

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate AD in samples. An antibody specific for AD has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyAD present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for AD is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of AD bound in the initial step. The color development is stopped and the intensity of the color is measured.

**Product Overview:**Adropin, a recently described endogenous neuroendocrine peptide, has been suggested to play a critical role in modulating normal physiological processes and maintaining metabolic homeostasis. Adropin (10 ng/mL) caused a marked 10-fold increase in endothelial cell migration and proliferation and significantly increased capillary-like endothelial cell sprouting and tube formation (p

Note: This product is for in vitro research use only