CD70 Antibody

Catalog No: #43184



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			400	
	esc	rın	tic	ı'n
\boldsymbol{L}	しつし	IIIU	uu	и I

CD70 Antibody	
Rabbit	
Polyclonal	
Antigen affinity purification.	
IHC	
Hu	
The antibody detects endogenous levels of total CD70 protein.	
peptide	
Synthetic peptide of human CD70	
CD70	
CD27L; CD27LG; TNFSF7	
Swiss-Prot#: P32970Gene ID: 970	
P32970	
970;	
2.4mg/ml	
Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.	
Store at -20°C	


Application Details

Immunohistochemistry: 1:40-1:250

Images



Immunohistochemical analysis of paraffin-embedded Human liver cancer tissue using #43184 at dilution 1/35.



Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using #43184 at dilution 1/35.

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin sythesis.

Note: This product is for in vitro research use only