

Ezrin (Phospho-Tyr478) Conjugated Antibody

Catalog No: #C11908

Package Size: #C11908-AF350 100ul #C11908-AF405 100ul #C11908-AF488 100ul

#C11908-AF555 100ul #C11908-AF594 100ul #C11908-AF647 100ul

#C11908-AF680 100ul #C11908-AF750 100ul #C11908-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | Ezrin (Phospho-Tyr478) Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of Ezrin only when phosphorylated at tyrosine 478. |
| Immunogen Description | Peptide sequence around phosphorylation site of tyrosine 478 (P-V-Y(p)-E-P) derived from Human Ezrin. |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Cytovillin;EZRI;p81;VIL2;Villin 2 |
| Accession No. | Swiss-Prot#:P15311NCBI Gene ID:7430NCBI mRNA#:NM_001111077.1 NCBI Protein#: NP_001104547.1 |
| Uniprot | P15311 |
| GeneID | 7430; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 70 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Probably involved in connections of major cytoskeletal structures to the plasma membrane. In epithelial cells, required for the formation of microvilli and membrane ruffles on the apical pole. Along with PLEKHG6, required for normal macropinocytosis.

Note: This product is for in vitro research use only