

ZNF76 Conjugated Antibody

Catalog No: #C43789



Package Size: #C43789-AF350 100ul #C43789-AF405 100ul #C43789-AF488 100ul

#C43789-AF555 100ul #C43789-AF594 100ul #C43789-AF647 100ul

#C43789-AF680 100ul #C43789-AF750 100ul #C43789-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	ZNF76 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous levels of total ZNF76 protein.
Immunogen Description	Synthetic peptide of human ZNF76
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	ZNF523;Zfp523;D6S229E
Accession No.	Swiss-Prot#:P36508NCBI Gene ID:7629NCBI mRNA#:NCBI Protein#:NP_003418
Uniprot	P36508
GeneID	7629;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	62
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

ZNF76, also known as ZNF523 or Zfp523, is a transcriptional repressor expressed in the testis. It is the human homolog of the Xenopus Staf protein (selenocysteine tRNA gene transcription-activating factor) known to regulate the genes encoding small nuclear RNA and selenocysteine tRNA. ZNF76 localizes to the nucleus and exerts an inhibitory function on p53-mediated transactivation. ZNF76 specifically targets TFIID (TATA-binding protein). The interaction with TFIID occurs through both its N and C termini. The transcriptional repression activity of ZNF76 is predominantly regulated by lysine modifications, acetylation and sumoylation. ZNF76 is sumoylated by PIAS 1 and is acetylated by p300. Acetylation leads to the loss of sumoylation and a weakened TFIID interaction. ZNF76 can be deacetylated by HDAC1. In addition to lysine modifications, ZNF76 activity is also controlled by splice variants. Two isoforms exist due to alternative splicing. These isoforms vary in their ability to interact with TFIID.

Note: This product is for in vitro research use only