MLC-2 (Phospho-Thr17+Ser18) Antibody

Catalog No: #12404

Package Size: #12404-1 50ul #12404-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|-----------------------|---|
| Product Name | MLC-2 (Phospho-Thr17+Ser18) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. |
| Applications | WB |
| Species Reactivity | Hu Ms Rt |
| Specificity | MLC-2 (Phospho-Thr17+Ser18) Antibody detects endogenous levels of MLC-2 only when phosphorylated at |
| | Thr17+Ser18 |
| Immunogen Type | Peptide |
| Immunogen Description | A synthesized peptide derived from human MLC-2 (Phospho-Thr17+Ser18) |
| Target Name | MLC-2 |
| Modification | Phospho |
| Other Names | MLC-2, MYL12A, C-type mannose receptor 2, CLEC13E, CD280, CD280 antigen, Endocytic receptor 180, |
| | ENDO180, KIAA0709, MRC2, Mannose receptor, C type 2, UPARAP, Macrophage mannose receptor 2, |
| | UPAR-associated protein |
| Accession No. | Swiss-Prot#: P19105/O14950/P24844NCBI Gene ID: 10627/103910/10398 |
| Uniprot | P19105 |
| GeneID | 10627; |
| Target Species | human |
| Calculated MW | 18kd |
| Concentration | 1.0mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide |
| | and 50% glycerol. |
| Storage | Store at -20°C |

Application Details

Western blotting: 1:1000

Note: This product is for in vitro research use only