

Myosin Light Chain 2 (Phospho-Ser19) Conjugated Antibody

Catalog No: #C11114

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Package Size: #C11114-AF350 100ul #C11114-AF405 100ul #C11114-AF488 100ul

#C11114-AF555 100ul #C11114-AF594 100ul #C11114-AF647 100ul

#C11114-AF680 100ul #C11114-AF750 100ul #C11114-Biotin 100ul

Description

Product Name	Myosin Light Chain 2 (Phospho-Ser19) Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of Myosin Light Chain 2 only when phosphorylated at serine 19.
Immunogen Description	Peptide sequence around phosphorylation site of serine 19 (A-T-S(p)-N-V) derived from Human Myosin Light Chain 2.
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	LC20; MLC2;MRLC1;MYRL2;MLC-2C
Accession No.	Swiss-Prot#:P19105NCBI Gene ID:10398NCBI mRNA#:NM_006097.4NCBI Protein#:NP_006088.2
Uniprot	P19105
GeneID	10627;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	18
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Product Description

Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Background

Myosin regulatory subunit that plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis, receptor capping, and cell locomotion

Note: This product is for in vitro research use only