

PAK2 (Ab-141) Conjugated Antibody

Catalog No: #C33243

Package Size: #C33243-AF350 100ul #C33243-AF405 100ul #C33243-AF488 100ul

#C33243-AF555 100ul #C33243-AF594 100ul #C33243-AF647 100ul

#C33243-AF680 100ul #C33243-AF750 100ul #C33243-Biotin 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

| | |
|-----------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name | PAK2 (Ab-141) Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total PAK2 protein. |
| Immunogen Description | Synthesized non-phosphopeptide derived from human PAK2 around the phosphorylation site of serine 141 (Y-L-S(p)-F-T). |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | EC 2.7.11.1;gamma-PAK;kinase PAK2;P21-activated kinase 2;p21-activated kinase 2 |
| Accession No. | Swiss-Prot#:Q13177NCBI Gene ID:5062 |
| Uniprot | Q13177 |
| GeneID | 5062; |
| Excitation Emission | AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm |
| Calculated MW | 58 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250

AF405 conjugated: most applications: 1: 50 - 1: 250

AF488 conjugated: most applications: 1: 50 - 1: 250

AF555 conjugated: most applications: 1: 50 - 1: 250

AF594 conjugated: most applications: 1: 50 - 1: 250

AF647 conjugated: most applications: 1: 50 - 1: 250

AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Product Description

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Background

Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell motility, cell cycle progression, apoptosis or proliferation. Acts as downstream effector of the small GTPases CDC42 and RAC1. Activation by the binding of active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues. Full-length PAK2 stimulates cell survival and cell growth. Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration. Phosphorylates JUN and plays an important role in EGF-induced cell proliferation. Phosphorylates many other substrates including histone H4 to promote assembly of H3.3 and H4 into nucleosomes, BAD, ribosomal protein S6, or MBP. Additionally, associates with ARHGEF7 and GIT1 to perform kinase-independent functions such as spindle orientation control during mitosis. On the other hand, apoptotic stimuli such as DNA damage lead to caspase-mediated cleavage of PAK2, generating PAK-2p34, an active p34 fragment that translocates to the nucleus and promotes cellular apoptosis involving the JNK signaling pathway. Caspase-activated PAK2 phosphorylates MNK1 and reduces cellular translation.

Note: This product is for in vitro research use only