NLRP4 Conjugated Antibody

Catalog No: #C37751



Package Size: #C37751-AF350 100ul #C37751-AF405 100ul #C37751-AF488 100ul #C37751-AF555 100ul #C37751-AF594 100ul #C37751-AF647 100ul #C37751-AF680 100ul #C37751-AF750 100ul #C37751-Biotin 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Beeenpaen | |
|-----------------------|--|
| Product Name | NLRP4 Conjugated Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Ни |
| Specificity | The antibody detects endogenous levels of total NLRP4 protein. |
| Immunogen Description | Synthetic peptide corresponding to a region derived from internal residues of human NLR family, pyrin domain |
| | containing 4 |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | CT58; PAN2; RNH2; NALP4; PYPAF4; CLR19.5 |
| Accession No. | Swiss-Prot#:Q96MN2NCBI Gene ID:147945NCBI Protein#:NP_653288 |
| Uniprot | Q96MN2 |
| GeneID | 147945; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |
| | |

Application Details

Suggested Dilution:

| AF350 conjugated: most applications: 1: 50 - 1: 250 |
|---|
| AF405 conjugated: most applications: 1: 50 - 1: 250 |
| AF488 conjugated: most applications: 1: 50 - 1: 250 |
| AF555 conjugated: most applications: 1: 50 - 1: 250 |
| AF594 conjugated: most applications: 1: 50 - 1: 250 |
| AF647 conjugated: most applications: 1: 50 - 1: 250 |
| AF680 conjugated: most applications: 1: 50 - 1: 250 |
| AF750 conjugated: most applications: 1: 50 - 1: 250 |
| Biotin conjugated: working with enzyme-conjugated str |

Background

NALPs are cytoplasmic proteins that form a subfamily within the larger CATERPILLER protein family. Most short NALPs, such as NALP4, have an N-terminal pyrin (MEFV; MIM 608107) domain (PYD), followed by a NACHT domain, a NACHT-associated domain (NAD), and a C-terminal leucine-rich repeat (LRR) region. The long NALP, NALP1 (MIM 606636), also has a C-terminal extension containing a function to find domain (FIIND) and a caspase recruitment domain (CARD). NALPs are implicated in the activation of proinflammatory caspases (e.g., CASP1; MIM 147678) via their involvement in multiprotein complexes called inflammasomes

Note: This product is for in vitro research use only