Product Datasheet

ASK1(Phospho-Ser83) Antibody

Catalog No: #11178

Package Size: #11178-1 50ul #11178-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	ASK1(Phospho-Ser83) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of ASK1 only when phosphorylated at serine 83.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 83 (G-S-S(p)-V-G) derived from Human ASK1.
Target Name	ASK1
Modification	Phospho
Other Names	ASK-1; M3K5; MAP3K5; MAPK/ERK kinase kinase 5; MAPKKK5
Accession No.	Swiss-Prot: Q99683NCBI Protein: NP_005914.1
Uniprot	Q99683
GeneID	4217;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.

Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 155kd
Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Images


Storage



Western blot analysis of extracts from K562 cells using ASK1(Phospho-Ser83) Antibody #11178(Lane 2) and the same antibody preincubated with blocking peptide(Lane1).



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using ASK1(Phospho-Ser83) Antibody #11178(left) or the same antibody preincubated with blocking peptide(right).



Western blot analysis of extracts from 3T3 cells, treated with Anisomycin or calf intestinal phosphatase (CIP), using ASK1 (Phospho-Ser83) Antibody #11178.

Background

Component of a protein kinase signal transduction cascade. Phosphorylates and activates MAP2K4 and MAP2K6, which in turn activate the JNK and p38 MAP kinases, respectively. Overexpression induces apoptotic cell death.

Mabuchi S, et al. (2004) Endocrinology. 145(1): 49-58.

Yuan ZQ, et al. (2003) J Biol Chem. 278(26): 23432-23440.

Kim AH, et al. (2001) Mol Cell Biol. 21(3): 893-901.

Note: This product is for in vitro research use only