NADH-cytochrome b5 reductase 3 Polyclonal Conjugated Antibody

Catalog No: #C42369

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com



#C42369-AF555 100ul #C42369-AF594 100ul #C42369-AF647 100ul

#C42369-AF680 100ul #C42369-AF750 100ul #C42369-Biotin 100ul

Description

| Product Name | NADH-cytochrome b5 reductase 3 Polyclonal Conjugated Antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of total NADH-cytochrome b5 reductase 3 polyclonal antibody. |
| Immunogen Description | Recombinant human NADH-cytochrome b5 reductase 3 protein |
| Conjugates | Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750 |
| Other Names | Diaphorase-1,CYB5R3, DIA1 |
| Accession No. | Swiss-Prot#:P00387 |
| Uniprot | P00387 |
| GeneID | 1727; |
| Excitation Emission | AF350: 346nm/442nm |
| | AF405: 401nm/421nm |
| | AF488: 493nm/519nm |
| | AF555: 555nm/565nm |
| | AF594: 591nm/614nm |
| | AF647: 651nm/667nm |
| | AF680: 679nm/702nm |
| | AF750: 749nm/775nm |
| Calculated MW | 33 |
| Formulation | 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide |
| Storage | Store at 4°C in dark for 6 months |

Application Details

Suggested Dilution:

AF350 conjugated: most applications: 1: 50 - 1: 250 AF405 conjugated: most applications: 1: 50 - 1: 250 AF488 conjugated: most applications: 1: 50 - 1: 250 AF555 conjugated: most applications: 1: 50 - 1: 250 AF594 conjugated: most applications: 1: 50 - 1: 250 AF647 conjugated: most applications: 1: 50 - 1: 250 AF680 conjugated: most applications: 1: 50 - 1: 250 AF750 conjugated: most applications: 1: 50 - 1: 250

Background

DIAPH1 binds to GTP-bound form of Rho and to profilin. Acts in a Rho-dependent manner to recruit profilin to the membrane, where it promotes actin polymerization. It is required for cytokinesis, stress fiber formation, and transcriptional activation of the serum response factor. DFR proteins couple Rho and Src tyrosine kinase during signaling and the regulation of actin dynamics (By similarity). In hearing it may play a role in the regulation of actin polymerization in hair cells.

Note: This product is for in vitro research use only