# Acetyl-Histone H3-K36 pAb

Catalog No: #29711

Package Size: #29711-1 50ul #29711-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


### Description

Product Name	Acetyl-Histone H3-K36 pAb
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	WB,IF
Species Reactivity	Human,Mouse,Rat
Immunogen Description	A synthetic acetylated peptide around K36 of human Histone H3 (NP_003484.1).
Other Names	HIST3H3; H3.4; H3/g; H3FT; H3t; histone H3.1t
Accession No.	Swiss-Prot#:Q16695NCBI Gene ID:8290
Uniprot	Q16695
GeneID	8290;
Calculated MW	Refer to figures
Formulation	Avoid freeze / thaw cycles. Buffer: PBS with 50% glycerol, pH7.4.
Storage	Store at -20°C


### **Application Details**

WB 1:500 - 1:2000IF 1:50 - 1:100


### **Images**



Western blot analysis of extracts of various cell lines, using Acetyl-Histone H3-K36 at 1:1000 dilution.HeLa cells were treated by TSA (1 uM) at 37°C for 18 hours.NIH/3T3 cells were treated by TSA (1 uM) at 37°C for 18 hours.



Immunofluorescence analysis of C6 cells using Acetyl-Histone H3-K36 at dilution of 1:100.C6 cells were treated by TSA (1 uM) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K36 at dilution of 1:100.NIH/3T3 cells were treated by TSA (1 uM) at 37°C for 18 hours. Blue: DAPI for nuclear staining.

## Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Note: This product is for in vitro research use only