

Acetyl-Histone H3-K4 pAb

Catalog No: #29712



Package Size: #29712-1 50ul #29712-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Acetyl-Histone H3-K4 pAb
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	WB,IF
Species Reactivity	Human,Mouse,Rat
Immunogen Description	A synthetic acetylated peptide around K4 of human Histone H3 (NP_003484.1).
Other Names	HIST3H3; H3.4; H3/g; H3FT; H3t; histone H3.1t
Accession No.	Swiss-Prot#:Q16695NCBI Gene ID:8290
Uniprot	Q16695
GeneID	8290;
Calculated MW	Refer to figures
Formulation	Avoid freeze / thaw cycles. Buffer: PBS with 50% glycerol, pH7.4.
Storage	Store at -20°C


Application Details

WB1:500 - 1:2000IF1:50 - 1:100


Images




Immunofluorescence analysis of C6 cells using Acetyl-Histone H3-K4 at dilution of 1:100.C6 cells were treated by TSA (1 uM) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using Acetyl-Histone H3-K4 at dilution of 1:100. HeLa cells were treated by TSA (1 μ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using Acetyl-Histone H3-K4 at dilution of 1:100. NIH/3T3 cells were treated by TSA (1 μ M) at 37°C for 18 hours. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using Acetyl-Histone H3-K4 at 1:1000 dilution. NIH/3T3 cells were treated by TSA (1 μ M) at 37°C for 18 hours. C6 cells were treated by TSA (1 μ M) at 37°C for 18 hours.

Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is located separately from the other H3 genes that are in the histone gene cluster on chromosome 6p22-p21.3.

Note: This product is for in vitro research use only