NR1H2 Polyclonal Antibody

Catalog No: #29799

Package Size: #29799-1 50ul #29799-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _ | | | | | |
|---|----|-----|---|-----|-----|
| П | es | cri | n | łi، | ٦n |
| u | てつ | UH | v | w | ווע |

| Product Name | NR1H2 Polyclonal Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Isotype | IgG |
| Purification | Affinity purification |
| Applications | WB,IF |
| Species Reactivity | Human,Mouse,Rat |
| Immunogen Description | A synthetic peptide of human NR1H2 (NP_009052.3). |
| Other Names | NR1H2;LXR-b;LXRB;NER;NER-I;RIP15;UNR |
| Accession No. | Uniprot:P55055GeneID:7376 |
| Uniprot | P55055 |
| GeneID | 7376 |
| Calculated MW | 51kDa |
| SDS-PAGE MW | 51kDa |
| Formulation | PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| Storage | Store at -20°C. Avoid freeze / thaw cycles. |
| | |

Application Details

WB 1:500 - 1:2000IF 1:50 - 1:200

Images



Western blot analysis of extracts of various cell lines, using NR1H2 antibody.

Background

The liver X receptors, LXRA (NR1H3; MIM 602423) and LXRB, form a subfamily of the nuclear receptor superfamily and are key regulators of macrophage function, controlling transcriptional programs involved in lipid homeostasis and inflammation. The inducible LXRA is highly expressed in liver, adrenal gland, intestine, adipose tissue, macrophages, lung, and kidney, whereas LXRB is ubiquitously expressed. Ligand-activated LXRs form

obligate heterodimers with retinoid X receptors (RXRs; see MIM 180245) and regulate expression of target genes containing LXR response elements (summary by Korf et al., 2009 [PubMed 19436111]).[supplied by OMIM, Jan 2010]

Note: This product is for in vitro research use only