## PML(isoform 5) Polyclonal Antibody

Catalog No: #30348

Package Size: #30348-1 50ul #30348-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| _                 |       |     |     |
|-------------------|-------|-----|-----|
|                   | escri | nti | n   |
| $\boldsymbol{ u}$ | COUL  | μu  | ULI |

| Product Name          | PML(isoform 5) Polyclonal Antibody                                |
|-----------------------|---|
| Host Species          | Rabbit  |
| Clonality             | Polyclonal  |
| Isotype               | IgG   |
| Purification          | Affinity purification   |
| Applications          | WB  |
| Species Reactivity    | Human,Mouse,Rat   |
| Immunogen Description | Recombinant protein of human PML.                                 |
| Other Names           | PML; MYL; PP8675; RNF71; TRIM19; protein PML                      |
| Accession No.         | Swiss-Prot#:P29590NCBI Gene ID:5371                               |
| Uniprot               | P29590  |
| GeneID                | 5371;   |
| Calculated MW         | Refer to figures  |
| Formulation           | Avoid freeze / thaw cycles. Buffer: PBS with 50% glycerol, pH7.4. |
| Storage               | Store at -20°C  |

## **Application Details**

WB 1:500 - 1:2000

## **Images**



Western blot analysis of extracts of various cell lines, using PML at 1:1000 dilution.

## Background

The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to oncogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Extensive alternative splicing of this gene

| results in several variations of the protein's central and C-terminal regions; all variants encode the same N-terminus. Alternatively spliced transcript |
|--|
| variants encoding different isoforms have been identified.   |

Note: This product is for in vitro research use only