

MAPK6 Conjugated Antibody

Catalog No: #C28907



Package Size: #C28907-AF350 100ul #C28907-AF405 100ul #C28907-AF488 100ul
#C28907-AF555 100ul #C28907-AF594 100ul #C28907-AF647 100ul
#C28907-AF680 100ul #C28907-AF750 100ul #C28907-Biotin 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

Description

Product Name	MAPK6 Conjugated Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	most applications
Species Reactivity	Hu,Ms,Rt
Immunogen Description	Recombinant fusion protein of human MAPK6 (NP_002739.1).
Conjugates	Biotin AF350 AF405 AF488 AF555 AF594 AF647 AF680 AF750
Other Names	MAPK6; ERK3; HsT17250; PRKM6; p97MAPK; mitogen-activated protein kinase 6
Accession No.	Swiss-Prot#:Q16659NCBI Gene ID:5597
Uniprot	Q16659
GeneID	5597;
Excitation Emission	AF350: 346nm/442nm AF405: 401nm/421nm AF488: 493nm/519nm AF555: 555nm/565nm AF594: 591nm/614nm AF647: 651nm/667nm AF680: 679nm/702nm AF750: 749nm/775nm
Calculated MW	120kDa
Formulation	0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6, 5mg/ml Bovine Serum Albumin, 0.02% Sodium Azide
Storage	Store at 4°C in dark for 6 months

Application Details

Suggested Dilution:

- AF350 conjugated: most applications: 1: 50 - 1: 250
- AF405 conjugated: most applications: 1: 50 - 1: 250
- AF488 conjugated: most applications: 1: 50 - 1: 250
- AF555 conjugated: most applications: 1: 50 - 1: 250
- AF594 conjugated: most applications: 1: 50 - 1: 250
- AF647 conjugated: most applications: 1: 50 - 1: 250
- AF680 conjugated: most applications: 1: 50 - 1: 250

AF750 conjugated: most applications: 1: 50 - 1: 250

Biotin conjugated: working with enzyme-conjugated streptavidin, most applications: 1: 50 - 1: 1,000

Background

The protein encoded by this gene is a member of the Ser/Thr protein kinase family, and is most closely related to mitogen-activated protein kinases (MAP kinases). MAP kinases also known as extracellular signal-regulated kinases (ERKs), are activated through protein phosphorylation cascades and act as integration points for multiple biochemical signals. This kinase is localized in the nucleus, and has been reported to be activated in fibroblasts upon treatment with serum or phorbol esters.

Note: This product is for in vitro research use only